

Concise Review: New Insights Into the Role of Macrophages in β -Cell Proliferation

XIANGWEI XIAO, GEORGE K. GITTES

Key Words. β -Cell proliferation • Macrophages • β -Cell regeneration • Diabetes • Macrophage polarization

Division of Pediatric Surgery,
Children's Hospital of
Pittsburgh, University of
Pittsburgh School of
Medicine, Pittsburgh,
Pennsylvania, USA

Correspondence: Xiangwei Xiao,
M.D., Ph.D., Division of Pediatric
Surgery, Children's Hospital of
Pittsburgh, University of
Pittsburgh School of Medicine,
4401 Penn Avenue, Pittsburgh,
Pennsylvania 15224, USA.
Telephone: 412-692-7290;
E-Mail: Xiangwei.xiao@chp.edu;
or George K. Gittes, M.D., Division
of Pediatric Surgery, Children's
Hospital of Pittsburgh, University
of Pittsburgh School of Medicine,
4401 Penn Avenue, Pittsburgh,
Pennsylvania 15224, USA.
Telephone: 412-692-7290;
E-Mail: gittesgk@upmc.edu

Received November 3, 2014;
accepted for publication March 9,
2015.

©AlphaMed Press
1066-5099/2015/\$20.00/0

[http://dx.doi.org/
10.5966/sctm.2014-0248](http://dx.doi.org/10.5966/sctm.2014-0248)

ABSTRACT

Diabetes mellitus can potentially be treated with islet transplantation, but additional sources of β cells are necessary to overcome the short supply of donor pancreases. Although controversy still exists, it is generally believed that the postnatal expansion of the β -cell mass is mainly through pre-existing β -cell replication. Thus, understanding the molecular mechanisms underlying the regulation of β -cell proliferation might lead to clinical strategies for increasing β -cell numbers, both in vitro and in vivo. Macrophages have a well-recognized role in the development of insulinitis as part of the pathogenesis of type 1 diabetes. However, a potential role for macrophage polarization, triggered by specific environmental stimuli, in promoting β -cell proliferation has only recently been appreciated. In the present review, we discuss several independent studies, using different regeneration models, that demonstrate a substantial inductive role for macrophages in β -cell proliferation. Additional dissection of the involved cell-cell crosstalk through specific signal transduction pathways is expected to improve our understanding of β -cell proliferation and might facilitate the current β -cell replacement therapy. *STEM CELLS TRANSLATIONAL MEDICINE* 2015;4:1–4

SIGNIFICANCE

New independent findings from different β -cell regeneration models, contributed by different research groups, have provided compelling evidence to highlight a previously unappreciated role for macrophages in β -cell proliferation. Additional dissection of the underlying mechanisms and cell-cell crosstalk might shed new light on strategies to increase the functional β -cell mass in vivo and on β -cell replacement therapies.

INTRODUCTION

Diabetes mellitus can potentially be treated with islet transplantation. Nevertheless, the short supply of donor pancreases constitutes a formidable obstacle to widespread clinical applications [1–5]. Although great efforts have been made to identify, isolate, and purify β -cell progenitors in the adult pancreas [6–13], mounting evidence suggests that β -cell neogenesis does not significantly contribute to the functional β -cell mass in the adult pancreas [14–28]. Most reports of β -cell transdifferentiation required genetic manipulations [29, 30].

Thus, increasing attention has been drawn to the induction of β -cell replication in vitro and in vivo, because mature β cells have a very slow proliferation rate [4] that declines further with age [31–36]. Macrophages have a well-recognized role in the development of insulinitis as part of the pathogenesis of type 1 diabetes (T1D). However, a potential role for macrophage polarization, triggered by specific environmental stimuli, in promoting

β -cell proliferation has only recently been appreciated [37–40]. In the present review, we discuss several independent studies using different regeneration models to demonstrate a substantial role for macrophages in β -cell proliferation. Additional dissection of the involved cell-cell crosstalk through specific signal transduction pathways is expected to improve our understanding of β -cell proliferation and might facilitate the current β -cell replacement therapy.

MACROPHAGE BIOLOGY

Macrophages are a type of white blood cell that engulf and digest cellular debris, foreign substances, microbes, and cancer cells in a process called phagocytosis. In addition to the macrophages that display this classic phenotype, designated M1 macrophages, another macrophage subtype, designated M2, is entirely different. The degree to which a given macrophage bears M1 or M2 characteristics is termed “polarization.” The M1 macrophages are induced by T helper 1 (Th1)

cell-derived interferon- γ and microbial products and respond to microbial infection with an enhanced phagocytic capability through the expression of inducible nitric oxide synthase and the secretion of proinflammatory cytokines, such as tumor necrosis factor- α , interleukin-12 (IL-12), IL-1 β , and IL-23, and toxic mediators, such as reactive oxygen species and nitric oxide [41–44]. M2 macrophages are induced during Th2-type responses through stimulation with IL-4/IL-13 and are responsible for wound healing and tissue remodeling functions [41–44]. Specifically, M2 macrophages are known to secrete a wide range of chemokines, enzymes, and growth factors to promote neovascularization, fibrosis, and tissue repair [41–44]. The M1/M2 polarization of tissue-destructive versus tissue-reparative macrophages is now regarded as a simplified categorization of a complex cell lineage, and more and more macrophage phenotypes in between the two polar opposites are being described [41–44].

MACROPHAGES IN T1D AND TYPE 2 DIABETES

It has been shown that islets do contain resident macrophages [45, 46]. Macrophages play an essential role in the development of T1D. The inactivation of macrophages in nonobese diabetic (NOD) mice prevents the occurrence of diabetes [47], because an environment with macrophages is necessary for differentiation of anti- β cell-cytotoxic T cells [48]. M1 macrophages, along with a subpopulation of CD4+ T lymphocytes that secrete high levels of IL-17 (Th17 cells) and CD8+ cytotoxic T cells, are together considered to be the major cell types that promote the development of T1D [49–52].

Obesity induces insulin resistance, which is a predisposing factor for the development of type 2 diabetes (T2D). Insulin resistance is promoted by a transition in macrophage polarization from the M2 activation state, maintained by STAT6 and peroxisome proliferator-activated receptors, to a classic M1 activation state driven by nuclear factor- κ B (NF- κ B), activator protein 1, and other related factors [53–56]. Previous studies in T2D have shown that M1 macrophages are associated with increasing inflammation, obesity, and insulin resistance, and M2 macrophages are associated with a reduction in both obesity and insulin resistance [49, 50]. Thus, macrophages play a nonredundant role in the pathogenesis of both T1D and T2D [57].

Very recently, epigenetic modifications in spontaneous autoimmune diseases have been identified. In line with this concept, a small-molecule inhibitor of a family of bromodomain-containing transcriptional regulators was recently shown to induce macrophages to adopt an anti-inflammatory phenotype, which suppressed the development of T1D in NOD mice in an NF- κ B-dependent manner [58]. In this model, islet inflammation was inhibited, and β cell regeneration occurred [58].

MACROPHAGES IN β -CELL PROLIFERATION

Macrophages, especially M2 macrophages, have been shown to secrete a variety of trophic factors, including vascular endothelial growth factor (VEGF), epidermal growth factor (EGF), transforming growth factor- β (TGF- β), angiopoietins, and Wnts, to regulate development, tissue remodeling, and tissue repair. However, evidence of their role in promoting β -cell proliferation has only been reported recently [37–40].




Figure 1. Signaling pathways through which macrophages regulate β -cell replication. M2 macrophages, not only release TGF- β 1 to directly induce upregulation of SMAD7 in β cells, but also release EGF to activate EGFR signaling that inhibits TGF- β 1-activated SMAD2 nuclear translocation, resulting in inhibition of TGF- β receptor signaling. SMAD7 promotes β -cell proliferation by increasing CyclinD1 and CyclinD2 and by inducing nuclear exclusion of p27. M2 macrophages also secrete Wnt ligands to activate the Wnt signaling pathway, which induces nuclear translocation and retention of β -catenin to promote β -cell replication. Abbreviations: EGF, epidermal growth factor; EGFR, EGF receptor; TGF- β 1, transforming growth factor- β 1.

First, in a regulatable β cell-specific VEGF overexpression mouse model, which results in significant β cell loss, withdrawal of VEGF overexpression led to β -cell proliferation, replenishing the reduced β -cell mass [39]. In this model, β -cell proliferation was found to depend on the recruitment of bone marrow-derived macrophages and their subsequent crosstalk with islet endothelial cells. Trophic factors released locally by recruited macrophages were proposed as a trigger for β -cell proliferation [39].

In another β cell regeneration model, in which mesenchymal stem cells were transplanted into the streptozotocin-treated mouse pancreas, significant β -cell proliferation was detected, contributing to the regeneration of functional β -cell mass [40]. In this model, recruited macrophages were found to be required for the β -cell proliferation to occur. In addition, Wnt signaling was proposed to be responsible for the cross-talk between M2 macrophages and β cells to induce β -cell proliferation [40] (Fig. 1).

In a β -cell injury/regeneration model, triggered by diphtheria toxin receptor-mediated conditional targeted cell death [59], β -cell regeneration was found to be attributable to the proliferation of surviving β cells, which was dependent on recruited M2 macrophages, involving the Wnt signaling pathway [38] (Fig. 1).

In addition to these three independent studies, we used another model of enhanced β -cell proliferation, partial pancreatic ductal ligation (PDL) [37]. We have previously compared two experimental models of β -cell proliferation, partial pancreatectomy (PPX) and PDL [19, 20, 60]. β -Cell proliferation after PPX is robust, presumably resulting from an increased workload demand on the residual β cells after surgical removal of a significant amount of the functional β -cell mass. In contrast, inflammation appears to be the major trigger for β -cell proliferation after PDL [20]. In line with this view, tissue injury in the PDL pancreas is accompanied by substantial infiltration of inflammatory cells, which appears to create a proliferation niche for β cells, such as has been seen in other regeneration models [38–40]. With the help of a recently

developed intraductal infusion system [18, 61, 62], we found that M2 macrophages, not only release TGF- β 1 to directly induce upregulation of SMAD7 in β -cells, but also release EGF to activate EGF receptor signaling, which inhibits TGF- β 1-stimulated SMAD2 nuclear translocation, resulting in inhibition of TGF- β receptor signaling. SMAD7 promotes β -cell proliferation by increasing CyclinD1 and CyclinD2 and by inducing nuclear exclusion of p27 [37] (Fig. 1).

FUTURE DIRECTIONS

From these recent studies, we hypothesize that a proliferation niche is needed for efficient β -cell proliferation and that macrophages might play a pivotal role in establishing this niche [37–40, 63]. β -Cell proliferation might involve the coordination of multiple processes, such as the detachment of cell-cell contacts, modulation of extracellular matrix, and release of growth factors, both locally and systemically [64]. Macrophages, with their high plasticity and phenotypic diversity, appear to play a critical role in β -cell proliferation by creating crosstalk among different cell types, including β cells, non- β endocrine cells, endothelial cells, mesenchymal cells, and other circulation-derived blood cells (Fig. 2). Additional elucidation of this crosstalk might substantially enhance our understanding of β -cell proliferation.

Because acinar cells have been shown to reprogram into insulin-producing cells by soluble growth factors [13] or genetic alterations [29, 30], it would be interesting to examine the role of macrophages in these models. Moreover, although a role for M2 macrophages in promoting β -cell proliferation in mice has been well described, any similar role for M2 macrophages in human β -cell proliferation has not been examined. A coculture experiment could potentially be applied to address this point.

CONCLUSION

Taken together, new independent findings from different β -cell regeneration models, contributed by different research groups, have provided compelling evidence to highlight a previously unappreciated role for macrophages in β -cell proliferation. Additional dissection of the underlying mechanisms and cell-cell crosstalk could shed new light on strategies to increase the functional β -cell mass in vivo and on β -cell replacement therapies.

REFERENCES

- 1 Pipeleers D, Keymeulen B, Chatenoud L et al. A view on beta cell transplantation in diabetes. *Ann N Y Acad Sci* 2002;958:69–76.
- 2 Zaret KS, Grompe M. Generation and regeneration of cells of the liver and pancreas. *Science* 2008;322:1490–1494.
- 3 Weir GC, Bonner-Weir S. Islet transplantation as a treatment for diabetes. *J Am Optom Assoc* 1998;69:727–732.
- 4 Ackermann AM, Gannon M. Molecular regulation of pancreatic beta-cell mass development, maintenance, and expansion. *J Mol Endocrinol* 2007;38:193–206.
- 5 Pipeleers D, Chintinne M, Denys B et al. Restoring a functional beta-cell mass in diabetes. *Diabetes Obes Metab* 2008;10(suppl 4):S4–62.

- 6 Kushner JA, Weir GC, Bonner-Weir S. Ductal origin hypothesis of pancreatic regeneration under attack. *Cell Metab* 2010;11:2–3.
- 7 Kopp JL, Dubois CL, Hao E et al. Progenitor cell domains in the developing and adult pancreas. *Cell Cycle* 2011;10:1921–1927.
- 8 Thorel F, Nèpote V, Avril I et al. Conversion of adult pancreatic alpha-cells to beta-cells after extreme beta-cell loss. *Nature* 2010;464:1149–1154.
- 9 Chera S, Baronnier D, Ghila L et al. Diabetes recovery by age-dependent conversion of pancreatic δ -cells into insulin producers. *Nature* 2014;514:503–507.
- 10 Chung CH, Hao E, Piran R et al. Pancreatic β -cell neogenesis by direct conversion from mature α -cells. *STEM CELLS* 2010;28:1630–1638.
- 11 Hao E, Lee SH, Levine F. Efficient β -cell regeneration by a combination of neogenesis and




Figure 2. M2 macrophages play an essential role in creating a proliferation niche for β -cell proliferation. β -Cell proliferation could involve the coordination of multiple processes, such as detachment of cell-cell contacts, modulation of extracellular matrix, and release of growth factors, both locally and systemically. Macrophages, with their high plasticity and phenotypic diversity, appear to play a critical role in β -cell proliferation by creating crosstalk among different cell types, including β cells, non- β endocrine cells, endothelial cells, mesenchymal cells, and other circulation-derived blood cells.

ACKNOWLEDGMENTS

This work was supported by NIH Grant R01 DK098196 (to G.K.G.), the Cochrane-Weber Endowed Fund in Diabetes Research (Grant NO19831 to X.X.), and the Children's Hospital of Pittsburgh Foundation.

AUTHOR CONTRIBUTIONS

X.X. and G.K.G.: manuscript writing, final approval of manuscript.

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

The authors indicated no potential conflicts of interest.

does not involve specialized progenitors. *Dev Cell* 2007;12:817–826.

16 Meier JJ, Butler AE, Saisho Y et al. Beta-cell replication is the primary mechanism subserving the postnatal expansion of beta-cell mass in humans. *Diabetes* 2008;57:1584–1594.

17 Georgia S, Bhushan A. Beta cell replication is the primary mechanism for maintaining postnatal beta cell mass. *J Clin Invest* 2004;114:963–968.

18 Xiao X, Guo P, Shiota C et al. Neurogenin3 activation is not sufficient to direct duct-to-beta cell transdifferentiation in the adult pancreas. *J Biol Chem* 2013;288:25297–25308.

19 Xiao X, Chen Z, Shiota C et al. No evidence for β cell neogenesis in murine adult pancreas. *J Clin Invest* 2013;123:2207–2217.

20 Xiao X, Wiersch J, El-Gohary Y et al. TGF β receptor signaling is essential for inflammation-induced but not β -cell workload-induced β -cell proliferation. *Diabetes* 2013;62:1217–1226.

21 Rankin MM, Wilbur CJ, Rak K et al. β -Cells are not generated in pancreatic duct ligation-induced injury in adult mice. *Diabetes* 2013;62:1634–1645.

22 Solar M, Cardalda C, Houbracken I et al. Pancreatic exocrine duct cells give rise to insulin-producing beta cells during embryogenesis but not after birth. *Dev Cell* 2009;17:849–860.

23 Kopp JL, Dubois CL, Schaffer AE et al. Sox9+ ductal cells are multipotent progenitors throughout development but do not produce new endocrine cells in the normal or injured adult pancreas. *Development* 2011;138:653–665.

24 Chintinne M, Stangé G, Denys B et al. Beta cell count instead of beta cell mass to assess and localize growth in beta cell population following pancreatic duct ligation in mice. *PLoS One* 2012;7:e43959.

25 Kopinke D, Brailsford M, Shea JE et al. Lineage tracing reveals the dynamic contribution of Hes1+ cells to the developing and adult pancreas. *Development* 2011;138:431–441.

26 Desai BM, Oliver-Krasinski J, De Leon DD et al. Preexisting pancreatic acinar cells contribute to acinar cell, but not islet beta cell, regeneration. *J Clin Invest* 2007;117:971–977.

27 Pan FC, Bankaitis ED, Boyer D et al. Spatiotemporal patterns of multipotentiality in Ptf1a-expressing cells during pancreas organogenesis and injury-induced facultative restoration. *Development* 2013;140:751–764.

28 Cavelti-Weder C, Shtessel M, Reuss JE et al. Pancreatic duct ligation after almost complete β -cell loss: Exocrine regeneration but no evidence of β -cell regeneration. *Endocrinology* 2013;154:4493–4502.

29 Zhou Q, Brown J, Kanarek A et al. In vivo reprogramming of adult pancreatic exocrine cells to beta-cells. *Nature* 2008;455:627–632.

30 Li W, Nakanishi M, Zumsteg A et al. In vivo reprogramming of pancreatic acinar cells to three islet endocrine subtypes. *Elife* 2014;3:e01846.

31 Hija A, Salpeter S, Klochendler A et al. G0-G1 transition and the restriction point in pancreatic β -cells in vivo. *Diabetes* 2014;63:578–584.

32 Salpeter SJ, Khalailah A, Weinberg-Corem N et al. Systemic regulation of the age-related

decline of pancreatic β -cell replication. *Diabetes* 2013;62:2843–2848.

33 Rankin MM, Kushner JA. Adaptive beta-cell proliferation is severely restricted with advanced age. *Diabetes* 2009;58:1365–1372.

34 Gunasekaran U, Gannon M. Type 2 diabetes and the aging pancreatic beta cell. *Aging (Albany, NY Online)* 2011;3:565–575.

35 Tonne JM, Sakuma T, Munoz-Gomez M et al. Beta cell regeneration after single-round immunological destruction in a mouse model. *Diabetologia* 2015;58:313–323.

36 Blum B, Roose AN, Barrandon O et al. Reversal of β cell de-differentiation by a small molecule inhibitor of the TGF β pathway. *Elife* 2014;3:e02809.

37 Xiao X, Gaffar I, Guo P et al. M2 macrophages promote beta-cell proliferation by up-regulation of SMAD7. *Proc Natl Acad Sci USA* 2014;111:E1211–E1220.

38 Criscimanna A, Coudriet GM, Gittes GK et al. Activated macrophages create lineage-specific microenvironments for pancreatic acinar- and β -cell regeneration in mice. *Gastroenterology* 2014;147:1106–1118.e11.

39 Brissova M, Aamodt K, Brahmachary P et al. Islet microenvironment, modulated by vascular endothelial growth factor-A signaling, promotes β cell regeneration. *Cell Metab* 2014;19:498–511.

40 Cao X, Han ZB, Zhao H et al. Transplantation of mesenchymal stem cells recruits trophic macrophages to induce pancreatic beta cell regeneration in diabetic mice. *Int J Biochem Cell Biol* 2014;53:372–379.

41 Flavell RA, Sanjabi S, Wrzesinski SH et al. The polarization of immune cells in the tumour environment by TGF β . *Nat Rev Immunol* 2010;10:554–567.

42 Gordon S, Taylor PR. Monocyte and macrophage heterogeneity. *Nat Rev Immunol* 2005;5:953–964.

43 Pollard JW. Trophic macrophages in development and disease. *Nat Rev Immunol* 2009;9:259–270.

44 Grunewald M, Avraham I, Dor Y et al. VEGF-induced adult neovascularization: Recruitment, retention, and role of accessory cells. *Cell* 2006;124:175–189.

45 Westwell-Roper CY, Ehses JA, Verchere CB. Resident macrophages mediate islet amyloid polypeptide-induced islet IL-1 β production and β -cell dysfunction. *Diabetes* 2014;63:1698–1711.

46 Calderon B, Suri A, Miller MJ et al. Dendritic cells in islets of Langerhans constitutively present beta cell-derived peptides bound to their class II MHC molecules. *Proc Natl Acad Sci USA* 2008;105:6121–6126.

47 Lee KU, Kim MK, Amano K et al. Preferential infiltration of macrophages during early stages of insulinitis in diabetes-prone BB rats. *Diabetes* 1988;37:1053–1058.

48 Jun HS, Yoon CS, Zbytniuk L et al. The role of macrophages in T cell-mediated autoimmune diabetes in nonobese diabetic mice. *J Exp Med* 1999;189:347–358.

49 Eguchi K, Manabe I. Macrophages and islet inflammation in type 2 diabetes. *Diabetes Obes Metab* 2013;15(suppl 3):152–158.

50 Espinoza-Jimenez A, Peon AN, Terrazas LI. Alternatively activated macrophages in types 1 and 2 diabetes. *Mediators Inflamm* 2012;2012:815953.

51 Honkanen J, Nieminen JK, Gao R et al. IL-17 immunity in human type 1 diabetes. *J Immunol* 2010;185:1959–1967.

52 Boehm BO, Rosinger S, Sauer G et al. Protease-resistant human GAD-derived altered peptide ligands decrease TNF-alpha and IL-17 production in peripheral blood cells from patients with type 1 diabetes mellitus. *Mol Immunol* 2009;46:2576–2584.

53 Jourdan T, Godlewski G, Cinar R et al. Activation of the Nlrp3 inflammasome in infiltrating macrophages by endocannabinoids mediates beta cell loss in type 2 diabetes. *Nat Med* 2013;19:1132–1140.

54 Han MS, Jung DY, Morel C et al. JNK expression by macrophages promotes obesity-induced insulin resistance and inflammation. *Science* 2013;339:218–222.

55 Menghini R, Casagrande V, Menini S et al. TIMP3 overexpression in macrophages protects from insulin resistance, adipose inflammation, and nonalcoholic fatty liver disease in mice. *Diabetes* 2012;61:454–462.

56 Koliwad SK, Streeper RS, Monetti M et al. DGAT1-dependent triacylglycerol storage by macrophages protects mice from diet-induced insulin resistance and inflammation. *J Clin Invest* 2010;120:756–767.

57 Olefsky JM, Glass CK. Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol* 2010;72:219–246.

58 Fu W, Farache J, Clardy SM et al. Epigenetic modulation of type-1 diabetes via a dual effect on pancreatic macrophages and β cells. *Elife* 2014;3:e04631.

59 Criscimanna A, Speicher JA, Houshmand G et al. Duct cells contribute to regeneration of endocrine and acinar cells following pancreatic damage in adult mice. *Gastroenterology* 2011;141:1451–1462.

60 El-Gohary Y, Tulachan S, Wiersch J et al. A SMAD signaling network regulates islet cell proliferation. *Diabetes* 2014;63:224–236.

61 Xiao X, Prasad K, Guo P et al. Pancreatic duct cells as a source of VEGF in mice. *Diabetologia* 2014;57:991–1000.

62 Xiao X, Guo P, Prasad K et al. Pancreatic cell tracing, lineage tagging and targeted genetic manipulations in multiple cell types using pancreatic ductal infusion of adeno-associated viral vectors and/or cell-tagging dyes. *Nat Protoc* 2014;9:2719–2724.

63 Van Gassen N, Van Overmeire E, Leuckx G et al. Macrophage dynamics are regulated by local macrophage proliferation and monocyte recruitment in injured pancreas. *Eur J Immunol* 2015 [Epub ahead of print].

64 Nikolova G, Jabs N, Konstantinova I et al. The vascular basement membrane: A niche for insulin gene expression and Beta cell proliferation. *Dev Cell* 2006;10:397–405.

Concise Review: New Insights Into the Role of Macrophages in β -Cell Proliferation

Xiangwei Xiao and George K. Gittes

Stem Cells Trans Med published online April 21, 2015

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stemcellstm.alphamedpress.org/content/early/2015/04/21/sctm.2014-0248>